

MULTIPLICITIES AND SPECTRA OF PROMPT FISSION NEUTRONS UP TO 200 MeV

Vladimir M. Maslov¹, Nikolai V. Kornilov², Alexander B. Kagalenko², Natalia A. Tetereva¹

¹ *Joint Institute for Nuclear and Energy Research -Sosny, 220109, Minsk-Sosny, Belarus*

² *Institute of Physics and Power Engineering, Obninsk, Russia*

The energy distributions of prompt fission neutrons for E_n up to 200 MeV are of great importance for the accelerator-driven system technologies of sustainable energy production and radioactive waste transmutation. Model calculations were performed to interpret the prompt fission neutron spectra (PFNS) of $^{232}\text{Th}(n,f)$ and $^{238}\text{U}(n,f)$ reaction for $E_n \lesssim 20$ MeV recently [1]. Pre-fission $^{238}\text{U}(n,xnf)$ reaction neutron spectra were calculated with a Hauser-Feshbach statistical model, ^{238}U fission and (n,xn) reaction cross section data being fitted. The average energy of the pre-fission (n,nf) neutrons is shown to be rather dependent on E_n . For $E_n = 6 - 9$ MeV the lowering of PFNS average energy, which is due to the pre-fission (n,nf) neutrons, is reproduced. For $E_n = 13 - 18$ MeV a lowering of the measured PFNS average neutron energies was interpreted. Spectra of neutrons, evaporated from fission fragments, were approximated as a sum of two Watt' distributions. Recently energy distributions of prompt fission neutrons for incident neutron energies up to 200 MeV were reported [2], multiplicity data for $^{235}\text{U}(n,f)$ and $^{238}\text{U}(n,f)$ might appear soon [3]. We would extend our phenomenological approach up to 200 MeV. A self-consistent model on the basis of the energy balance with the incorporation of chance structure of fission would be employed [4]. It was verified below 20 MeV. The calculated values are compatible also with experimental data for $E_n \sim 20-50$ MeV for the ^{232}Th , ^{235}U , ^{238}U target nuclides. We will explore the energy dependence of the pre- and post-fission neutron multiplicities. Realistic calculations of average energies of pre-fission neutrons would be carried out for E_n up to 200 MeV for the consistent neutron multiplicity analysis at high incident energies for (n,f) and (p,f) reactions. Importance of separate measurements for the ν_{pre} and ν_{post} for (n,f) and (p,f) reactions would be exemplified.

References

- [1] V.M. Maslov et al., EuroPhysics Journal A, 18 (2003) 93.
- [2] Ethvignot T. et al., Phys. Lett. B. (2003), in press.
- [3] Ethvignot T., private communication (2003).
- [4] Kornilov N.V., Kagalenko A.B., Maslov V.M., et al, preprint IPPE-2978, 2003.